

Pregnancy Risk Factors as found in WHO and other developing world sources (December 1994)

As analysed and sometimes edited by

Rupert Fawdry, FRCS (Ed), FRCOG.

Consultant Specialist in Obstetrics, Gynaecology and Medical Information Technology
General Editor: Electronic Encyclopaedia of Perinatal Data (EPPD) Web site: www.fawdry.info
31, St.Mary's Way, Leighton Buzzard, LU7 2RX, United Kingdom
Home: 01525 37 01 37 Fax: 01525 38 24 53 Mobile: 077 68 97 44 13
e-mail: eepd@fawdry.demon.co.uk

or

c/o The Perinatal Institute, Crystal Court,
Aston Cross (off Rocky Lane), Birmingham B6 5RQ
Tel: 0121 687 3400 www.perinatal.nhs.uk

**Any Comments, Criticisms, Corrections
or Suggestions for Improvement very welcome**

LATE PREGNANCY CHECK.(LAST TWO MONTHS).

GENERAL.

No previous antenatal care.

Pallor.

Anaemia. Hb less than 10 gm/l.

Poor weight gain.

Weight gain 5kg or less at 24 weeks gestation.

6kg or less at the end of pregnancy.

Total weight gain less than 42kg at 34 weeks gestation.

Blood pressure 90 or more diastolic.

Albuminuria.

Glycosuria

High fasting blood sugar.

High blood sugar 2 hours after a normal meal

(Post prandial blood sugar).

ABDOMEN.

Malpresentation.

Fundal height too small.

too big.

(using coloured tape)

VAGINAL

Vaginal Bleeding.

Clinical assessment of pelvis.

FETUS.

Movements decreased.